

张学忠采。

分布：中国（四川）。

5 皱背无齿股蟊 *Parasipylodea rugulosa* Chen et He, 2008

Parasipylodea rugulosa Chen et He, 2008: 166, fig. 132. Type locality: Menglong, Yunnan, China.

研究标本：未见标本。

分布：中国（云南）。

致谢 承河北大学博物馆借用标本，在此特致谢意。

REFERENCES

- Chen, S-C and He, Y-H 1994. Newly recorded genus and a new species of Heteronemiidae (Phasmatodea). *Journal of Beijing Forestry University*, 16 (4): 121 – 123.
- Chen, S-C and He, Y-H 2008. Phasmatodea of China. China Forestry Publishing House, Beijing. 1 – 476.
- Ho, G. W. C. 2013. A new genus and five new species of Necrosiinae (Phasmatodea, Diapheromeridae) from China. *Acta Zootaxonomica Sinica*, 38 (3): 530 – 534. [动物分类学报]
- Ho, G. W. C. 2013. New records of Chinese Necrosiinae (Phasmatodea: Diapheromeridae). *Hong Kong Entomological Bulletin*, 5 (1): 1 – 5.
- Redtenbacher, J. 1908. Die Insektenfamilie der Phasmiden. III. Phasmidae Anareolatae (Phibalosomini, Acrophyllini, Necrosiini). Wilhelm Engelmann, Leipzig. 341 – 589.
- Vickery, V. R. 1983. Catalogue of Australian stick insects (Phasmida, Phasmatodea, Phasmatoptera, or Cheleutoptera). *CSIRO Australian division of Entomology, Technical Paper*, 20: 1 – 19.